

ACTIVITY SHEET: ONE-SAMPLE T-TEST FOR POPULATION MEAN

This activity sheet includes exercises to assess students' understanding of important concepts presented in the *One-Sample t-Test for Population Mean* lesson.

One-Sample t -Test for Population Mean

The data for these exercises are in the Minitab file **OneSampleTTest_PopMean_Activity.mtw**.

Exercise 1

For the following multiple-choice and true/false problems, choose the correct answer.

(a) True or False. The t values that corresponds to $\alpha = 0.01$ in both tails of the t distribution with 5 degrees of freedom are $\pm t_{0.005,5} = \pm 4.032$.

(b) True or False. The t distribution has "heavier" tails than the normal distribution, especially for small values of the sample size n . That is, there is more area in the tails of a t distribution for $n < 15$ than there is for the normal distribution.

(c) True or False. In a test of $H_0: \mu = 8$ versus $H_a: \mu \neq 8$, a one-sample t -test results in a p -value of 0.034. Thus, if we create a 95% two-sided confidence interval for μ with the data used to perform the hypothesis test, the interval will include $\mu = 8$.

(d) True or False. You are conducting the hypothesis test $H_0: \mu = 12$ versus $H_a: \mu \neq 12$ for a normally distributed population in which σ is unknown. For a sample size of $n = 10$, you obtain the standardized test statistic $t = 2.04$. The p -value for this hypothesis test is approximately 0.07 (rounded correctly to 2 decimal places).

(e) A horticulturist wants to estimate the mean growth of seedlings in a large timber plot planted last year. The growth of seedlings in these conditions is known to be normally distributed. A random sample of $n = 10$ seedlings is selected and the one-year growth for each is measured. The sample results are: $\bar{x} = 5.62$ cm and $s = 2.50$ cm. The 90% two-sided confidence interval for the mean growth μ is:

A. [5.16, 6.08] cm



- B. [4.17, 7.07] cm
- C. [3.12, 8.12] cm
- D. [4.98, 6.26] cm
- E. [4.32, 6.92] cm

(f) True or False. The average length of time that an engine is allowed to remain running after it is cranked ("run time") is a major factor influencing the longevity of the engine. A new model car is tuned such that it will experience optimal engine life if the true mean run time is 12.5 minutes. An owner of this new model car records her run time for all the trips that she drives during the third month of ownership. Below is a random sample of 12 of her run times.

12.6 8.6 12.2 25.4 46.4 36.5 9.8 9.7 19.9 9.5 8.7 25.7

To perform a hypothesis test to determine if the true mean run time μ for her car is 12.5 minutes, she should use a one-sample t -test with $df = 11$.

(g) True or False. As an entrepreneur, Lucy sets up a Psychiatric stand instead of the traditional Lemonade stand. She decides to collect some data to determine if, on average, she makes more than \$3.75 per day from her stand. She takes a systematic random sample of 15 days (ordered by day) over the summer and records the amount she makes each day. The correct test for her to perform to address her hypothesis is a **one-sample t -test**. Below is a summary of the data.



(h) Suppose we are constructing a two-sided confidence interval for the true population mean by using a simple random sample with sample size $n = 50$. What is the confidence level associated with the following confidence interval for μ ?

$$\bar{x} \pm 2.575 \cdot s/\sqrt{50}$$

- A. 99.5%

- B. 99.35%
- C. 99%
- D. 98.7%
- E. 98%

(i) An operations research analyst for a hotel agency has been asked to develop a fairly accurate estimate of the true mean "check-in" time for customers arriving in the hotel's lobby at noon. The estimate will be used to determine the number of front desk employees required so that a customer's wait time is "reasonable." Suppose the analyst randomly samples 16 customers' check-in times over the next month. She finds that their average wait time is 4.2 minutes with a standard deviation of 1.4 minutes. Which of the following is the correct two-sided 90% confidence interval for the true mean check-in time μ .

- A. (3.586, 4.814) minutes
- B. (3.454, 4.946) minutes
- C. (3.514, 4.886) minutes
- D. (3.624, 4.776) minutes
- E. We don't have enough information to construct this confidence interval.

(j) True or False. I talk to my mom on the phone every night. I randomly selected 11 dates (in minutes) up to July 4, 2020 and recorded the phone time usage with my mom. Here is the data:

13.34 35.23 45.12 32.14 33.82 75.23 15.22 42.22 35.68 40.24 33.21

The data is normally distributed at level of significance $\alpha = 0.05$.

(k) True or False. In conducting the hypothesis test $H_0: \mu = 8$ versus $H_a: \mu \neq 8$, the corresponding p -value is 0.44. If we construct a 95% two-sided confidence interval for μ using the exact same data, the value 8 would not be included in this 95% confidence interval.

- A. True
- B. False
- C. We do not have enough information to determine this.

(l) Suppose you perform the following hypothesis test on the true mean GPA at College A: $H_0: \mu = 3.1$ versus $H_a: \mu > 3.1$. You obtain a p -value for the hypothesis test. Which of the following statements is correct regarding the p -value?

- A. Under the assumption that H_0 is true, an extremely small p -value is consistent with the sample mean \bar{x} greatly differing from null mean ($\mu = 3.1$).
- B. The p -value measures the probability that the alternative hypothesis is true.
- C. The p -value measures the probability that the null hypothesis is true.
- D. The larger the p -value, the stronger the evidence against the null hypothesis.

E. A large p -value indicates that the data supports with the alternative hypothesis.

(m) Suppose a company boasts that it has new fertilizer that yields *at least* 2 tons (per year) of a certain crop over 50 acres. Twenty farmers (whose 50 acres usually yield 2 tons) are randomly selected to try the new fertilizer on their crops.

What are the null hypothesis, alternative hypothesis, test statistic and conclusion for testing the company's claim at the $\alpha = 0.05$ significance level? Use the Minitab output below and assume the population (crop growth) is approximately normally distributed.

Descriptive Statistics

N	Mean	StDev	SE Mean
20	1.9800	0.3200	0.0716

μ : population mean of crop growth

A. $H_0: \mu = 2$ versus $H_a: \mu > 2$, $t_0 = -0.28$. The company does not have enough evidence to reject the null hypothesis at $\alpha = 0.05$. Thus, their sample data *does not support* the new fertilizer yielding a larger crop.

B. $H_0: \mu = 2$ versus $H_a: \mu < 2$, $t_0 = -0.28$. The company does not have enough evidence to reject the null hypothesis at $\alpha = 0.05$. Thus, their sample data *does support* the new fertilizer yielding a larger crop.

C. $H_0: \mu = 2$ versus $H_a: \mu > 2$, $t_0 = -0.28$. The company does have enough evidence to reject the null hypothesis at $\alpha = 0.05$. Thus, their sample data *does support* that the new fertilizer is yielding a larger crop.

D. $H_0: \mu = 2$ versus $H_a: \mu < 2$, $z_0 = -0.28$. The company does not have enough evidence to reject the null hypothesis at $\alpha = 0.05$. Thus, their sample data *does not support* the new fertilizer yielding a larger crop.

E. $H_0: \mu = 2$ versus $H_a: \mu \neq 2$, $z_0 = -1.25$. The company does not have enough evidence to reject the null hypothesis at $\alpha = 0.05$. Thus, their sample data *does not support* the new fertilizer yielding a larger crop.

(n) What is the p -value associated with the following hypothesis test with sample data as shown below?

$$H_0: \mu = 28 \text{ vs } H_a: \mu \neq 28$$

Descriptive Statistics

N	Mean	StDev	SE Mean
42	28.500	1.630	0.252

μ : population mean

A. There is not enough information to determine an answer

B. 0.000

C. 0.054

- D. 0.023
- E. 0.027
- F. 0.047

(o) We are performing a one-sample t -test on a population mean μ in Minitab. Assume that we are told that the population is normally distributed. Here are the null and alternative hypotheses:

$$H_0: \mu = 75 \text{ vs } H_a: \mu \neq 75$$

Suppose we only obtain the following Minitab results. Use these results to answer the two questions below.

Descriptive Statistics

N	Mean	StDev	SE Mean	99% CI for μ
20	70.85	6.56	1.47	(66.65, 75.05)

μ : population mean of home prices in a certain city

- i. **True or False.** H_0 should be rejected at level of significance $\alpha = 0.05$.
- ii. **True or False.** H_0 should be rejected at level of significance $\alpha = 0.01$.

Exercise 2 based on the movie “Beauty and the Beast”

To win first prize at the inventor’s fair, Maurice designs a machine that improves wood chopping. He is interested in establishing that, on average, his machine can chop *more than* 420 lbs. of wood per hour. He conducts a study in which the machine is run for 20 one-hour intervals. At the end of each hour, he weighs the amount of wood chopped.

- (a) State the null and alternative hypothesis appropriate for addressing Maurice’s question of interest. Be sure to define symbols used to represent the parameter.
- (b) Using the data that he collected, Maurice computes a test statistic of $t_0 = 2.38$. Assuming all necessary assumptions are reasonable for using a t distribution to model his data (e.g. normality), compute the p -value that accompanies Maurice’s output.
- (c) Assuming that all necessary assumptions are reasonable, what conclusions can Maurice make at the $\alpha = 0.05$ significance level? Be sure to state the conclusions in the context of the problem.

Exercise 3 based on the movie “Beauty and the Beast”

As a gift to Belle, the Beast presented her with a library full of books. Belle is determined to read every book in the library. Whether she can accomplish her goal partly depends on the length of

the books. Let μ represent the average length (number of pages) of a book in the library. Belle is interested in determining if:

$$H_0: \mu = 500 \text{ vs. } H_a: \mu < 500$$

She takes a random sample of 50 books and records the number of pages contained in each. Below is a partial printout of the Minitab output corresponding to her analysis.

N	Mean	t
50	476.4	-1.04

μ : population mean of the number of book pages

(a) After looking at the results from Belle's analysis, Beast states that "there is insufficient evidence, at the $\alpha = 0.05$ level, to conclude that there are less than 500 pages in a book, on average." Is his statement justified? Explain.

(b) Which of the following **is** an appropriate interpretation of the p -value for this test?

- A. The probability that the alternative hypothesis is true is 0.151.
- B. The probability that the null hypothesis is true is 0.151.
- C. In repeated sampling, the probability of observing a test statistic less than or equal to -1.04 if the alternative hypothesis is true is 0.151.
- D. In repeated sampling, the probability of observing a test statistic less than or equal to -1.04 if the null hypothesis is true is 0.151.


(c) Which of the following would be an appropriate two-sided confidence interval to assess the hypothesis $H_0: \mu = 500$ vs. $H_a: \mu < 500$ at the $\alpha = 0.02$ significance level?

96% z-interval	98% z-interval	99% z-interval
96% t-interval	98% t-interval	99% t-interval

(d) Suppose she repeats the study increasing the sample size to $n = 100$. Holding all else fixed, what effect would this have on the reported p -value of 0.151?

- A. The p -value will decrease.
- B. The p -value will remain the same.
- C. The p -value will increase.
- D. It is not possible to say how the p -value will change.

(c) Lumiere and Cogsworth each are responsible for overseeing the castle. Suppose they are interested in testing whether Plumette, a feather duster, requires *more than* 8 hours, on average, to clean the entire castle. Cogsworth and Lumiere decide to collect separate samples, each taking a sample of size 20. A side-by-side boxplot of their data is shown below. Which character will obtain a *larger* test statistic for assessing the question of interest? Explain.



Exercise 4: Tootsie Pops

Below are data results for a random sample of $n = 13$ students in an Engineering Statistics class who were given Tootsie Pop suckers and asked to record how long it took (in seconds) until they first tasted the inside tootsie roll center.

205 440 425 192 195 225 260 465 306 450 385 460 468

Using the appropriate method, determine a two-sided 90% confidence interval for the true mean amount of time to get to the center of a Tootsie Pop for these students. Report your final confidence interval values correct to 2 decimal places.

Exercise 5: Restaurant Clean-up Times

A restaurant employee (with a statistics background) at "Sushi Yummy" has been asked to provide a 99% two-sided confidence interval the true mean time to clean off a restaurant table and get it prepared for the next customer on a busy Saturday night. This knowledge will help the restaurant owners know how much time they should schedule between reservations. Suppose the employee randomly samples table clean up times for $n = 40$ tables on busy Saturday nights and obtains a sample mean clean-up time of $\bar{x} = 4.2$ minutes with a sample standard deviation of $s = 0.8$ minutes.

The sample data is used to construct the following two-sided confidence interval for the true mean clean up time μ :

(3.962, 4.438) minutes

Determine the approximate confidence level (e.g., 75%, 88%, 95%) of this confidence interval correct to 2 decimal places.

Exercise 6: Low Tire Pressure

(From the article *Whatever Happened to...Full Service Gas Stations?* at Go Retro.^{*}) The oil crisis of the 1970s marked the beginning of the end for the full-service gas station. Oil companies figured that customers wanted to pump their own gas in exchange for saving a few pennies. Pretty soon, the attendants were no longer needed. Also, the process of getting gas at a full-service station took about 10-15 minutes, which sadly is considered too long in today's high paced, impatient world.

^{*} <https://www.goretro.com/2011/07/whatever-happened-tofull-service-gas.html>



As a result of gas stations moving from full-service to self-service, a certain gas station in a small town is concerned that many cars are being driven on underinflated tires. This can result in excessive tire wear, and unsafe steering and braking of the car. A tire is seriously underinflated if its tire pressure is more than 10 psi under its recommended level.

The gas station manager selects a random sample of 100 cars at his station and records the cars' front driver's side tire pressure while the customer pumps their gas. He determines that the mean underinflation for this random sample is 10.4 psi with a standard deviation of $s = 4.2$ psi.

- (a) Construct a 99% two-sided confidence interval for the mean underinflation μ .
- (b) Based on your confidence interval from part (a), would you recommend that the manager issue a report that the mean tire pressure is seriously underinflated? Explain your answer.

Exercise 8: Taking Fire

The World War I Flying Ace (a.k.a. Snoopy) often searches the skies for his enemy, The Red Baron. As a result, his aircraft often sustains damage from shots fired from the Baron. The Ace's commanding officers are interested in estimating the average number of bullet holes in the side of the Ace's aircraft after an encounter with the Baron. For a random sample of 14 flights in which the Ace encountered the Baron, the number of bullet holes in the side of the aircraft is recorded. The data for this problem is in the column "Number of Bullet Holes" in the Minitab worksheet associated with this lesson's activities.

(a) The commanding officers would like to construct a confidence interval for the mean number of bullets; they would like to use a “t-interval.” Based on the data for this problem, is this the correct interval to construct? Explain.

(b) Regardless of what you concluded in part (a), the officers constructed a 98% “t-interval” to estimate the mean and obtained the following interval: (12.61, 19.39) bullet holes. What critical value (cv) was used to construct this interval? See the expression below for further clarification on the value cv you are determining.

$$\bar{x} \pm cv \cdot \frac{s}{\sqrt{n}} \rightarrow (12.61, 19.39) \text{ bullet holes}$$

(c) Regardless of your answer to part (a), assume the underlying population is normally distributed. The Flying Ace states that he is able to escape from the Red Baron with fewer than 15 bullet holes, on average. Using the interval provided in part (b), is his statement justified? Explain.

(d) Regardless of your answer to part (a), assume the underlying population is normally distributed. Which of the following is an appropriate interpretation of the interval stated in part (b)?

- A. 98% of flights observed had between 12.61 and 19.39 bullet holes.
- B. 98% of all flights have between 12.61 and 19.39 bullet holes.
- C. There is a 98% chance that the average number of bullet holes for this sample is between 12.61 and 19.39.
- D. There is a 98% chance that the average number of bullet holes in the population is between 12.61 and 19.39.
- E. None of the above is a correct interpretation.

(e) Suppose the officers were to repeat the study collecting a new random sample of 14 flights and observed a sample standard deviation of 0. Which of the following statements *must* be true?

- A. Half of the values observed are above the sample mean.
- B. All of the values observed are zero.
- C. All of the values observed are equivalent.
- D. The values are evenly spaced on both sides of the sample mean.

